

What Does It Take to Stabilize Gravitational Clustering?

Chung-Pei Ma

*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia PA 19104;
cpma@physics.upenn.edu*

and

J. N. Fry

Department of Physics, University of Florida, Gainesville FL 32611-8440; fry@phys.ufl.edu

ABSTRACT

An analytical understanding of the strongly nonlinear regime of gravitational collapse has been difficult to achieve. The only insight has been the stable clustering hypothesis, which assumes that the number of neighbors for objects averaged over small length scales is constant in time. Our recently proposed analytic halo model for N -point correlation functions now provides a tool for calculating gravitational clustering properties in the strongly nonlinear regime. This model also provides a new physical framework for an independent evaluation of the validity of the stable clustering hypothesis. We derive the asymptotic nonlinear behavior of the N -point correlation functions and pairwise peculiar velocities in terms of dark matter halo properties. We show that these statistics exhibit stable clustering only when the halo mass function and halo density profile obey specific relations. The long-cherished stable clustering hypothesis therefore is not necessarily realized in practice.

Subject headings: cosmology : theory – dark matter – large-scale structure of universe

1. Introduction

Gravitational clustering is the fundamental physical process responsible for the formation and evolution of structure in the universe. Galaxy-hosting dark matter halos are the products of the strongly nonlinear stage of this process, but a detailed understanding of this important regime has mostly eluded us because few numerical simulations have the dynamic range to explore comfortably such a small length scale (e.g., Moore et al. 1999; Bullock et al. 1999). The one analytic handle has been the stable clustering assumption (Peebles 1974; Davis & Peebles 1977; Peebles 1980), which has allowed us to extrapolate into nonlinear regimes beyond the reach of numerical simulations. Predictions of the stable clustering hypothesis are: (1) the two-point correlation function of the density field $\xi(r)$ and the power spectrum $P(k)$ are power laws, with $\xi(r) \propto r^{-\gamma}$

and $\Delta(k) \equiv 4\pi k^3 P(k) \propto k^\gamma$, where $\gamma = (9 + 3n)/(5 + n)$ for a primordial spectral index n ; (2) the higher order N -point correlation functions $\xi_N(r)$ scale as $\xi_N(r) \propto r^{-\gamma_N}$, with $\gamma_N = (N - 1)\gamma$, or $\xi_N \propto \xi^{N-1}$; and (3) the pairwise peculiar velocity exactly cancels the Hubble expansion on small scales, $v/Hr = -1$, so that bound, high-density halos maintain a fixed physical size. Some of these predictions have agreed well enough with numerical simulation results (Jain 1997 and references therein) to have obtained general acceptance.

In two recent papers we presented evidence from high-resolution N -body simulations that gravitational clustering does not necessarily follow the scaling $\xi_3 \propto \xi^2$ required by the stability condition for the two- and three-point functions ξ and ξ_3 (Ma & Fry 2000a). For a deeper understanding of the strongly nonlinear regime beyond simulations, we proceeded to construct an analytic halo model (Ma & Fry 2000b) in which mass is distributed in spherical dark matter halos with phenomenological mass distribution functions, density profiles, and halo-halo correlations. We showed that this halo model can reproduce analytically the two- and three-point correlation functions measured in numerical simulations, and that it also makes useful predictions beyond the limited range of validity of simulations.

In this *Letter* we investigate the extent to which the analytic halo model is consistent with stable gravitational clustering in the strongly nonlinear regime. In particular, we derive the halo model predictions for the asymptotic nonlinear behavior of the N -point correlation functions (§2) and the pairwise peculiar velocities (§3) in terms of dark matter halo properties. We then obtain the conditions that must be satisfied by the halo mass function and density profile in order to reproduce results of the stable clustering hypothesis.

2. N -Point Correlation Functions

The analytic halo model of Ma & Fry (2000b) for the N -point correlation functions of the mass density takes three inputs: the halo mass distribution function dn/dM ; the halo density profile $\rho(r)/\bar{\rho} = \bar{\delta} u(r/R_s)$ where R_s is a scale radius and $\bar{\delta}$ is the normalization; and halo-halo correlations. In this model, the two-point function contains contributions from particle pairs residing in a single halo and in two separate halos, $\xi(r) = \xi_{1h}(r) + \xi_{2h}(r)$, where the subscripts “1h” and “2h” denote the respective “1-halo” and “2-halo” contributions (see also Seljak 2000). Similarly, we write the bispectrum $B(k_1, k_2, k_3)$ as three separate terms, representing contributions from particle triplets that reside in a single halo and in two and three distinct halos. As shown in Ma & Fry (2000b), the dominant contribution in the nonlinear regime on small length scales is from the “1-halo” terms for particles that reside in the same halos. This makes intuitive sense because closely spaced particles are most likely to be found in the same halo. These “1-halo” terms for the two- and N -point statistics are

$$\xi_{1h}(r) = \int dM \frac{dn}{dM} R_s^3 \bar{\delta}^2 \lambda_2(r/R_s), \quad P_{1h}(k) = \int dM \frac{dn}{dM} [R_s^3 \bar{\delta} \tilde{u}(kR_s)]^2,$$

$$\begin{aligned}\xi_{N,1h}(r_1, \dots, r_N) &= \int dM \frac{dn}{dM} R_s^3 \bar{\delta}^N \lambda_N(r_1/R_s, \dots, r_N/R_s), \\ P_{N,1h}(\mathbf{k}_1, \dots, \mathbf{k}_N) &= \int dM \frac{dn}{dM} [R_s^3 \bar{\delta} \tilde{u}(k_1 R_s)] \cdots [R_s^3 \bar{\delta} \tilde{u}(k_N R_s)].\end{aligned}\tag{1}$$

where $\lambda_N(x_1, \dots, x_N) \equiv \int d^3y u(y) u(|\mathbf{y} + \mathbf{x}_1 - \mathbf{x}_2|) \cdots u(|\mathbf{y} + \mathbf{x}_1 - \mathbf{x}_N|)$; $\tilde{u}$ is the Fourier transform of $u(x)$; the N -point P_N is defined for $\sum \mathbf{k}_i = 0$, and $P_3 = B$ is the bispectrum.

For our goal of studying the N -point functions in the strongly nonlinear regime, we need to consider only the halo mass function and halo density profile and not halo-halo correlations because the latter make negligible contributions on small length scales and are therefore irrelevant for stable clustering (see Ma & Fry 2000b for the full expressions for P and B). For the halo mass function, we write it as

$$\frac{dn}{dM} \propto M^{-2} \nu^\alpha e^{-\nu^2/2}, \quad \nu = \frac{\delta_c}{\sigma(M)},\tag{2}$$

where α parameterizes the uncertainty in the logarithmic slope at the low-mass end, $\delta_c \approx 1.68$ characterizes the linear overdensity at the onset of gravitational collapse, and $\sigma(M)$ is the linear rms mass fluctuations in spheres of radius R , where $M = 4\pi\bar{\rho}R^3/3$. Values in use for α vary from 1 (Press & Schechter 1974) to about 0.4 (Sheth & Tormen 1999; Jenkins et al. 2000). For the halo density profile, we assume a “universal” shape, $\rho/\bar{\rho} = \bar{\delta} u(r/R_s)$, where $u = 1/[x(1+x)^2]$ is proposed by Navarro et al. (1997), while $u(x) = 1/(x^{3/2} + x^3)$ is suggested by Moore et al. (1999). Both the scale radius R_s and the normalization $\bar{\delta}$ can be expressed in terms of a concentration parameter $c(M) \equiv R_{200}/R_s$, where R_{200} is the virial radius within which the average halo density is 200 times the mean density of the universe. Specifically, $R_s \propto M^{1/3}/c$, and $\bar{\delta} \propto c^3 g(c)$, where $g(c)$ is a logarithmic factor with $g = 1/[\ln(1+c) - c/(1+c)]$ for Navarro et al. (1997) and $g = 1/\ln(1+c^{3/2})$ for Moore et al. (1999). High resolution simulations of individual halos show a roughly power-law $c(M)$ with some, perhaps substantial, scatter (e.g., Cole & Lacey 1996; Tormen et al 1997; Navarro et al. 1997; Jing & Suto 2000), so we will write

$$c \equiv \frac{R_{200}}{R_s} \propto \left(\frac{M}{M^*} \right)^{-\beta_0},\tag{3}$$

where M^* is defined by $\sigma(M^*) = 1$, and β_0 is used to parameterize the uncertainty in the mass dependence, which has been found to be in the range $0 \lesssim \beta_0 \lesssim 1/2$.

We now derive analytically the small- r behavior of equation (1), in which the factors dn/dM , R_s , and $\bar{\delta}$ are all functions of the halo mass M . For small r or high k , the integral for $\xi_{1h}(r)$ or $P_{1h}(k)$ converges before the exponential cutoff in the mass function, and the dominant contribution to the integral comes from masses near the scale for which $r/R_s = 1$ or $kR_s = 1$. We then have $P(k) \approx P_{1h}(k) \propto \int dM \nu^\alpha \tilde{u}^2(kR_s) g^2(c)$ at high k . Changing variables to $y = kR_s \propto k M^{\beta_0+1/3}$, we obtain the asymptotic behavior

$$\xi(r) \propto r^{-\gamma}, \quad \Delta(k) \equiv 4\pi k^3 P(k) \propto k^\gamma, \quad \gamma = \frac{18\beta - \alpha(n+3)}{2(3\beta+1)},\tag{4}$$

where we have $\beta = \beta_0$ if the logarithmic factor $g(c)$ in $\bar{\delta}$ is ignored, or more accurately, we have $\beta \approx 0.8\beta_0$ if the effect of $g(c)$ is approximated by a power law. From equation (4), we see that in order to reproduce the two-point stable clustering result for a given spectral index n , the indices α and β must satisfy

$$\frac{18\beta - \alpha(n+3)}{2(3\beta+1)} = \frac{9+3n}{5+n}. \quad (5)$$

For $n = -2$, for example, this is satisfied if $\beta = 0.25$ or $\beta_0 \approx 0.31$ for $\alpha = 1$, and $\beta = 0.2$ or $\beta_0 \approx 0.25$ for $\alpha = 0.4$.

For the three-point function ξ_3 and the bispectrum B , the integrals in equation (1) converge if $\epsilon < 3/2$ for an inner halo density of $r^{-\epsilon}$. For ξ_3 , B , and the commonly defined three-point amplitude $Q \propto B/P^2$, we find

$$\xi_3(r) \propto r^{-\gamma_3}, \quad k^6 B(k) \propto k^{\gamma_3}, \quad Q(k) \propto k^{\gamma_3-2\gamma}, \quad \gamma_3 = \frac{36\beta - \alpha(n+3)}{2(3\beta+1)}. \quad (6)$$

Stable clustering requires a scale-independent $Q(k)$, and

$$\gamma_3 - 2\gamma = \frac{\alpha(n+3)}{2(3\beta+1)} = 0, \quad (7)$$

which holds only if $\alpha = 0$ or $n = -3$. Combining equations (5) and (7), we find that stable clustering requires $\alpha = 0$ and $\beta = (3+n)/6$. All work on halo mass functions performed thus far has reported a much larger value for α ; stable clustering therefore does not appear to be followed, which is consistent with the scale-dependent $Q(k)$ reported in Ma & Fry (2000a).

We illustrate these results by showing in Figure 1 the power spectrum and bispectrum computed from the analytic halo model for three sets of (α, β) for the $n = -2$ scale-free model. The values of α and β are all chosen to give the same slope $\gamma = 1$ for the nonlinear two-point function; the corresponding power spectra are therefore nearly identical at high k . The three-point Q , however, have very different high- k behavior for the three cases, and only the case $\alpha = 0$ produces an approximately scale-independent Q . (Q is not perfectly flat for $(\alpha, \beta) = (0, 1/6)$ because as mentioned above, the simple expressions in eqs. (5) and (7) are derived by approximating the logarithmic factor $g(c)$ in the halo profile as a power law.)

For the four-point and higher correlation functions, provided the integral is convergent without the exponential cutoff in the mass function, the same change of variable gives

$$\xi_N \propto r^{-\gamma_N}, \quad k^{3(N-1)} P_N \propto k^{\gamma_N}, \quad \gamma_N = \frac{18(N-1)\beta - \alpha(n+3)}{2(3\beta+1)}. \quad (8)$$

Stable clustering requires

$$\gamma_N - (N-1)\gamma = (N-2)\frac{\alpha(n+3)}{2(3\beta+1)} = 0, \quad (9)$$

which again is satisfied by $\alpha = 0$ or $n = -3$. So if the three-point correlation follows equation (7), stable clustering continues to hold for all orders for which the integral converges. At large k , we are

probing the inner cores of halos, where the inner profile goes as $r^{-\epsilon}$. The Fourier amplitude $\tilde{u}$ thus falls off as $\tilde{u}(k) \propto (kR_s)^{-(3-\epsilon)}$, and convergence requires $N[1-(3-\epsilon)(3\beta+1)/3]-1+\alpha(n+3)/6 < 0$. For $\epsilon = 1$ (Navarro et al.) and $n > -2$, this holds for all N , but for $\epsilon = \frac{3}{2}$ (Moore et al.) and $n = -2$, it holds only as far as $N = 3$.

3. Pairwise Peculiar Velocities

Besides the scaling properties of the N -point correlation functions discussed above, stable clustering also predicts that the mean peculiar velocity v between pairs of objects of separation r must cancel the Hubble expansion Hr on small scales so that bound halos maintain a fixed physical size. The velocity ratio v/Hr is related to the two-point correlation function ξ by the pair conservation equation (Peebles 1980)

$$-\frac{v}{Hr} = \frac{\bar{\xi}}{3(1+\xi)} \frac{\partial \ln \bar{\xi}}{\partial \ln a}, \quad (10)$$

where a is the expansion factor and $\bar{\xi}(x) \equiv 3x^{-3} \int_0^x dx' x'^2 \xi(x')$ is the mean two-point correlation function interior to x . Stable clustering requires $-v/Hr = 1$.

The analytic halo model of Ma & Fry (2000b) makes specific predictions for the behavior of v/Hr . As eq. (10) indicates, the main task is to compute the time dependence of ξ . Equation (1) shows that the time dependence of ξ enters through the rms density fluctuation σ in dn/dM of eq. (2), and the halo concentration parameter $c(M)$ of eq. (3). We obtain

$$\frac{\partial \xi_{1h}}{\partial \ln a} = \int dM [f(\nu^2 - \alpha) + (2\bar{\delta}' - 3 + \lambda') c'] \frac{dn}{dM} R_s^3 \bar{\delta}^2 \lambda(r/R_s), \quad (11)$$

where $f \equiv d \ln \sigma / d \ln a \approx \Omega^{0.6}$ is the standard growth rate of the density field, $\bar{\delta}' \equiv d \ln \bar{\delta} / d \ln c$, $\lambda' \equiv d \ln \lambda / d \ln x$, and $c' \equiv d \ln c / d \ln a$. Specifically, for the Navarro et al. profile, $\bar{\delta}(c) = (200 c^3 / 3) / [\ln(1+c) - c/(1+c)]$ and $\lambda(x) = 8\pi[(x^2 + 2x + 2) \ln(1+x)/(x^2 + 2x) - 1]/(x^3 + 2x^2)$; and similar expressions for the Moore et al. profile can be found in Ma & Fry (2000b). The term c' for the time dependence of the concentration parameter is somewhat uncertain. Numerical simulation results of Bullock et al. (1999) give a stronger redshift evolution, $c \propto a$, than Navarro et al. (1997), while more complicated forms have also been proposed (e.g., Cooray et al. 2000). We note that in equation (3), $c(M)$ has an implicit dependence on a through M^* which roughly agrees with Bullock et al. when $\beta = (3+n)/6$.

The prediction of the halo model for v/Hr can be calculated from equations (1) and (11). Before doing so, it is useful to derive first the asymptotic behavior of v/Hr in the deeply nonlinear regime at small r . On such small scales, $\xi \gg 1$, and equation (4) gives a power law $\xi \propto r^{-\gamma}$ and $\bar{\xi} = 3\xi/(3-\gamma)$. For a scale-free model, the self similarity solution gives $\partial \bar{\xi} / \partial \ln a = -2/(3+n) \partial \bar{\xi} / \partial \ln r$. We find that equation (10) then reaches

$$-\frac{v}{Hr} \xrightarrow{r \rightarrow 0} \frac{2}{n+3} \frac{18\beta - \alpha(n+3)}{6 + \alpha(n+3)}. \quad (12)$$

It can be easily checked that $-v/Hr$ approaches unity, as it should, for any pair (α, β) that satisfies the stable clustering condition of equation (5) for the two-point function.

Figure 2 shows our results for the general behavior of $-v/Hr$ calculated from the analytic halo model. An $n = -2$ scale-free power spectrum is used for illustration. Although ξ is dominated by the 1-halo term ξ_{1h} of equation (1) at $\xi \gtrsim 1$, which is the range of interest for this paper, we improve the accuracy at $\xi < 1$ by adding the linear theory $\xi_l \propto r^{-(3+n)} \propto r^{-1}$ to ξ_{1h} . As Figure 2 shows, the resulting v/Hr indeed follows the linear perturbative prediction at $\xi \lesssim 1$, and equation (12) at $\xi \gg 1$. We find $-v/Hr \rightarrow 1$ for the same pairs of (α, β) shown in Figure 1, which were chosen to illustrate the stable clustering scaling of the two-point function. However, $-v/Hr$ reaches a smaller value at $\xi \gg 1$ for the (α, β) suggested in the literature, e.g., $\alpha = 1$ or 0.4 for the mass function (Press & Schechter 1974; Sheth & Tormen 1997; Jenkins et al. 2000) and $\beta = 1/6$ (Cole & Lacey 1996; Navarro et al. 1997), implying that the pairwise peculiar velocities are too small to cancel exactly the Hubble expansion. As a consistency check, we have computed the derivative $\partial \bar{\xi} / \partial a$ using both eq. (11) and $[\bar{\xi}(a + \Delta a) - \bar{\xi}(a)] / \Delta a$. The two methods give identical results, but eq. (11) offers a more physical insight into the various contributing terms. For example, the broad peak in $-v/Hr$ for $1 \lesssim \xi \lesssim 100$ is the result of the term in ν^2 in eq. (11), which reflects that the integral on these scales is dominated by mass scales near M^* where $\sigma(M^*) = 1$.

4. Summary and Discussion

In this *Letter*, we have derived the asymptotic nonlinear behavior of the N -point correlation functions ξ_N (eqs. [4], [6], [8]) and the pairwise peculiar velocities $-v/Hr$ (eq. [12]) in the framework of the recently proposed analytic halo model of Ma & Fry (2000b). We have shown that their small scale behavior is consistent with the stable clustering hypothesis only if dark matter halos satisfy certain criteria. The two halo parameters whose variation we have explored are the logarithmic slope α of the halo mass distribution dn/dM at the low mass end in eq. (2) and the slope β_0 for the mass dependence of the halo concentration parameter $c(M)$ in eq. (3) (recall $\beta \approx 0.8\beta_0$). These two parameters are highlighted because results to date from numerical simulations have indicated a significant uncertainty, with $0.4 \lesssim \alpha \leq 1$ and $0 \lesssim \beta \lesssim 1/2$ being plausible values.

From the derived asymptotic nonlinear behavior, we have obtained analytically the relations that must be satisfied by α and β in order for stable clustering to occur. Equations (5), (7), and (9) summarize the results for the N -point correlation function ξ_N . These equations and Figure 1 indicate that although the two-point stable clustering condition alone is satisfied by an infinite set of (α, β) for a given n , the three- and higher-point stable clustering conditions, $\xi_N \propto \xi^{N-1}$, $N \geq 3$, imply α must be zero. Achieving stable clustering to all orders therefore requires stringent conditions: $\alpha = 0$ and $\beta = (3 + n)/6$. For non-scale-free models (e.g., cold dark matter), n is the index at the high- k end of the power spectrum.

Compared with the hierarchical model of Davis & Peebles (1977), results from this paper and

Ma & Fry (2000b) show that the analytic halo model makes more general and physical predictions for the behavior of the N -point correlation functions and the pairwise velocities. Imposing the stability condition $\alpha = 0$ and $\beta = (3 + n)/6$ in equations (4), (6), and (8), we indeed recover their results of $\xi \propto r^\gamma$ with $\gamma = (9 + 3n)/(5 + n)$ at small r and $\xi_N \propto \xi^{N-1}$. The condition $\beta = (3 + n)/6$ can be understood further by examining individual halos. Using $c \propto M_*^{\beta_0}$, $M_* \propto a^{6/(n+3)}$, and $\beta_0 \approx \beta$ up to a logarithmic factor, we see that $\beta = (3 + n)/6$ gives approximately $c \propto a(t)$, which is consistent with the simulation results in Bullock et al. (2000). The halo scale radius R_s is then approximately constant in time in physical coordinates, as expected for a stable system, and the physical density ρ of individual halos is indeed very close to constant in time, more so for very small halos that form early ($\nu \gg 1$, $c \gg 1$) than for those that form recently ($\nu \approx 1$, $c \approx 1$).

Yano & Gouda (1999) have found that for halo density $\rho \propto r^{-\epsilon}$ with $\epsilon \leq 3/2$ (such as Navarro et al. and Moore et al.), the velocity parameter v/Hr approaches 0 and the two-point correlation function ξ approaches a constant (i.e. $\gamma = 0$) in the nonlinear limit. Neither behavior is seen in our Figures because their derivation ignores the mass distribution function dn/dM and is therefore valid only for equal mass halos.

Our earlier paper (Ma & Fry 2000a) has already shown signs of departure from the stable clustering hypothesis in high resolution N -body simulations. In this paper, the specific values $\alpha = 0$ and $\beta = (3 + n)/6$ required for stable clustering provide additional evidence that this long-cherished hypothesis may not be applicable in all situations. In such cases, one consequence is that the frequently used fitting formulas for the nonlinear power spectrum (Hamilton et al 1991; Jain et al. 1995; Peacock & Dodds 1996; Ma 1998) will need modifications at high k , e.g., $k/k_{nl} \gtrsim 50$ for the $n = -2$ model and $k \gtrsim 20 h \text{ Mpc}^{-1}$ for cold dark matter models with a cosmological constant (see Figs. 3 and 4 of Ma & Fry 2000b). We believe the analytic halo model is a powerful tool that is providing new insight into the nonlinear regime of gravitational clustering, the most fundamental process in cosmology.

C.-P. M thanks Jim Peebles and Uros Seljak for useful discussions. She acknowledges support of an Alfred P. Sloan Foundation Fellowship, a Cottrell Scholars Award from the Research Corporation, a Penn Research Foundation Award, and NSF grant AST 9973461.

REFERENCES

- Bullock, J. S. et al. 1999, astro-ph/9908159; submitted to MNRAS
- Cole, S., & Lacey, C. 1996, MNRAS, 281, 716
- Cooray, A., Hu, W., & Miralda-Escude, J. 2000, astro-ph/0003205
- Davis, M., & Peebles, P. J. E. 1977, ApJS, 34, 425
- Hamilton, A. J. S., Matthews, A., Kumar, P., & Lu, E. 1991, ApJ, 374, L1
- Jain, B. 1997, MNRAS, 287, 687
- Jain, B., Mo, H. J., & White, S. D. M. 1995, MNRAS, 276, L25
- Jenkins, A., Frenk, C. S., White, S. D. M., Colberg, J. M., Cole, S., Evrard, A. E., & Yoshida, N. 2000, astro-ph/0005260
- Jing, Y. P., & Suto, Y. 2000, ApJ, 529, L69
- Ma, C.-P. 1998, ApJ, 508, L5
- Ma, C.-P., & Fry, J. N. 2000a, ApJ, 531, L87
- Ma, C.-P., & Fry, J. N. 2000b, ApJ, 543, in press, astro-ph/0003343
- Moore, B., Quinn, T., Governato, F., Stadel, J., & Lake, G. 1999, MNRAS, 310, 1147
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
- Peacock, J. A., & Dodds, S. J. 1996, MNRAS, 280, L1
- Peebles, P. J. E. 1974, ApJ, 189, L51
- Peebles, P. J. E. 1980, The Large-Scale Structure of the Universe (Princeton: Princeton Univ. Press)
- Press, W. H., & Schechter, P. 1974, ApJ, 187, 425
- Seljak, U. 2000, astro-ph/0001493
- Sheth, R. K., & Tormen, G. 1999, MNRAS, 308, 119
- Tormen, G., Bouchet, F., & White, S. D. ., 1997, MNRAS, 286, 865
- Yano, T., & Gouda, N. 1999, astro-ph/9906375




Fig. 1.— The power spectrum (upper panel) and three-point amplitude (lower panel) given by the analytic halo model of Ma & Fry (2000b) for the $n = -2$ scale-free model. Three sets of (α, β) are shown, where α and β parameterize the halo mass function and halo concentration as defined in eqs. (2) and (3). (Note β_0 in eq. [3] is related to β here by $\beta \approx 0.8\beta_0$; see text.) The Navarro et al. profile with $c(M_*) = 9$ is assumed. Dotted straight lines show the lowest order perturbative predictions. While different combinations of (α, β) give nearly identical $\Delta(k)$ which all agree with the two-point stable clustering result (see eq. [5]), the three-point $Q(k)$ has distinct high- k behavior (see eq. [6]), and stable clustering holds to all orders only if $\alpha = 0$ and $\beta = (3 + n)/6$.




Fig. 2.— Prediction of the analytic halo model of Ma & Fry (2000b) for the ratio of pairwise peculiar velocity and Hubble flow, $-v/Hr$, vs. the two-point correlation function ξ for the $n = -2$ model. The velocity ratio approaches a constant in the deeply nonlinear regime (see eq. [12]), but the value is not always unity as required by stable clustering. For example, $-v/Hr \rightarrow 1$ at $\xi \gg 1$ for $(\alpha, \beta) = (1, 1/4), (0.4, 1/5), (0, 1/6)$, but it reaches a smaller value for $\alpha = 0.4$ or 1 and $\beta = 1/6$, which are the range of values found in simulations. Dotted straight line shows the linear theory prediction, which is followed accurately at $\xi \lesssim 1$.